

Dietary regimen: ☐ Vegetarian ☐ Vegan

Riomnis

Clinical information form

Functional coprology

INTERNATIONAL DIVISION - Tel.: +33 (0)4 72 80 23 85 - Fax: +33 (0)4 72 80 73 56 - E-mail: serviceexport@eurofins-biomnis.com **PATIENT DETAILS** First name(s): Surname: Date of birth: Gender: F Age: years **ELASTASE REPORT** Shipment of refrigerated stool sample, minimum 20 g (or discrete stool fragment). **CLINICAL DETAILS** YES Known exocrine pancreatic insufficiency: If yes, what is etiological diagnosis:

alcoholism cystic fibrosis ☐ idiopathic, medicinal, obstructive IP **Enzyme replacement therapy:** YES \square NO CALPROTECTIN **REPORT** Shipment of refrigerated stool sample, minimum 20 g (or discrete stool fragment). **CLINICAL DETAILS** Inflammatory bowel disease (Crohn's, ulcerative colitis):

YES NO YES IBD treatment in progress: If yes, what is treatment: ☐ salicylate derivatives corticosteroids ☐ immunosuppressors anti-TNF 24-HOUR STOOL ANALYSIS, STEATORRHEA AND CREATORRHEA FORM FOR RECEIPT AND SHIPMENT OF STOOLS Reception of stools for 24, 48 or 75 hours in sufficiently large pots provided by the laboratory. • Grouping saddles of 48 or 72h in ONE SAME BAG AND ONE SINGLE FILE so as to constitute a single examination carried out from a pool of saddles of 48 or 72h. Pots may not be more than 2/3 filled. Essential minimum volume = 30 g. • Check the screwing of the lid for a good closure of the pot to avoid leakage of samples. • Be sure to collect stools produced at night. • Refrigerate the stools to limit fermentation phenomena. • Each pot is to be identified by the family name, forename, date of birth and date of collection of the stool samples. • It is essential to provide the lab with information on the period over which the stools were collected. NOTES • Diet enriched with butter (50g butter minimum daily) 3 days before collection and during the collection for the assay of steatorrhea and fecalogramme. Avoid consumption of oil-rich foods (peanuts, walnuts, hazelnuts, avocados) during the 3 days preceding the collection and during the collection. · Avoid taking medications that slow the passage of stools, coloscopy preparations, suppositories, laxatives, intestinal dressings and radiological examinations with agents that opacify the bowel contents in the 3 days before collection and during collection itself. **CLINICAL DETAILS Gastrointestinal surgical history**: ☐ Gastrectomy ☐ Colectomy **Digestive pathology:** \square Irritable bowel syndrome ☐ Celiac disease ☐ EPI Other: **Endocrine / neuroendocrine disease:** ☐ Diabetes ☐ Hyperthyroidism Other: If yes, please specify: Weight loss: Average no. of stools per day:

☐ Gluten-free

Treatment in progress (digestive purposes):

□ None □ Other: